arXiv:1004.0442v1 [cond-mat.stat-mech] 3 Apr 2010

SHORT-WAVE EXCITATIONS IN NON-LOCAL GROSS-PITAEVSKII MODEL

A.P. IVASHIN, Yu.M. POLUEKTOV

A.I. Akhiezer Institute for Theoretical Physics,
NSC "Kharkov Institute of Physics and Technology"
(1, Akademicheskaya Str., Kharkov UA-61108, Ukraine;
e-mail: ivashin@kipt.kharkov.ua,yuripoluektov@kipt.kharkov.ua)

UDC 532.591,538.94,539.196.3

©2009

It is shown, that a non-local form of the Gross-Pitaevskii equation allows to describe not only the long-wave excitations, but also the short-wave ones in the systems with Bose-condensate. At given parameter values, the excitation spectrum mimics the Landau spectrum of quasi-particle excitations in superfluid Helium with roton minimum. The excitation wavelength, at which the roton minimum exists, is close to the inter-particle interaction range. It is shown, that the existence domain of the spectrum with a roton minimum is reduced, if one accounts for an inter-particle attraction.

The Gross-Pitaevskii equation (GP) obtained in [1, 2, 3] is widely used in researches of superfluid Bose-systems of weakly interacting particles. This equation describes the particle dynamics in a Bose-condensate. In particular, it gives a framework to study the condensate oscillations and to obtain the elementary excitation spectra. The knowledge of the latter is necessary in order to obtain the thermodynamic functions of the system.

At a stage of derivation of the GP equation for the condensate wave function $\Phi\left(r,t\right)$, it is usually assumed that Φ varies slowly over the distances of order r_0, which is the range of the inter-atomic interaction potential $U\left(r\right)$. This condition allows to express the GP equation in form of a non-linear differential equation.

Unfortunately, in such a case, the imposed assumption restricts a consideration to the long-wave excitations in the Bose-condensate. Namely, it allows one to derive the Bogolyubov sound (long-wave) oscillation spectrum; nevertheless, a (physical) superfluid system can also have some well-defined long-lived short-wave excitations with the de Broglie wavelengths of the order of inter-particle distance (or interaction potential range). E.g., in the superfluid helium the rotons are the excitations of this type.

The purpose of this paper is to demonstrate that the non-local form of the GP equation describes the short-wave excitations in a system with Bose-condensate, as well as the long-wave ones, if the constraint on the of macroscopic wave function spacial change scale is released. At given values of the parameters the excitation spectrum mimics the Landau one for the quasi-particles in the superfluid Helium, which has the roton minimum. We study the influence of the inter-particle interaction on the dispersion curve in the short-wave domain. It is shown that the inter-particle attraction narrows down the existence domain of a stable spectrum. In particular, it reduces the range of parameters where a spectrum with roton minimum may exist.

NON-LOCAL FORM OF GROSS-PITAEVSKII EQUATION

A system of many Bose-particles in the second quantization picture with an account for the pairwise interactions is described by the Hamiltonian

$$\begin{aligned} H \;=\; & \int d\mathbf{r}\hat{\psi}^{+}\left(\mathbf{r}\right)\left(-\frac{\hbar^2}{2m}\Delta\right)\hat{\psi}\left(\mathbf{r}\right) \\ & +\frac{1}{2}\iint d\mathbf{r}d\mathbf{r}'\hat{\psi}^{+}\left(\mathbf{r}\right)\hat{\psi}^{+}\left(\mathbf{r}'\right)U\left(\mathbf{r}-\mathbf{r}'\right)\hat{\psi}\left(\mathbf{r}'\right)\hat{\psi}\left(\mathbf{r}\right). \end{aligned} \tag{1}$$

The equation of motion for the field operator in the Heisenberg picture reads

$$\begin{aligned} i\hbar\frac{\partial\hat{\psi}\left(\mathbf{r},t\right)}{\partial t} \;=\; & -\frac{\hbar^2}{2m}\Delta\hat{\psi}\left(\mathbf{r},t\right) \\ & +\psi\left(\mathbf{r},t\right)\int d\mathbf{r}'\,U\left(\mathbf{r}-\mathbf{r}'\right)\left|\hat{\psi}\left(\mathbf{r}',t\right)\right|^2, \end{aligned} \tag{2}$$

Due to a large number of particles in the condensate, one may consider the operator $\hat{\psi}(\mathbf{r},t)$ in eq. (2) as a non-operator function $\psi\left(\mathbf{r},t\right)$. Further, it is convenient to define a function $\Phi\left(\mathbf{r},t\right)$:

$$\psi\left(\mathbf{r},t\right)=\Phi\left(\mathbf{r},t\right)\exp\left(-i\mu\, t\right). \tag{3}$$

Then the equation reads

$$\begin{aligned} i\hbar\frac{\partial\Phi\left(r,t\right)}{\partial t} \;=\; & -\left(\frac{\hbar^2}{2m}\Delta+\mu\right)\Phi\left(r,t\right) \\ & +\Phi\left(r,t\right)\int dr'\,U\left(r-r'\right)\left|\Phi\left(r',t\right)\right|^2, \end{aligned} \tag{4}$$

where μ is the chemical potential which value is determined by the equilibrium of the system

$$\mu = \int d\mathbf{r}'\, U(\mathbf{r}-\mathbf{r}')\, |\Phi_0(\mathbf{r}')|^2, \tag{5}$$

with $\Phi_0(\mathbf{r})$ the equilibrium value of the condensate wave function. Taking into account that the equilibrium condensate density in absence of external fields $n_0 = |\Phi_0(\mathbf{r})|^2$ does not depend on the coordinate, one can express eq. (4) as

$$\begin{aligned} i\hbar\frac{\partial\Phi(\mathbf{r},t)}{\partial t} &= -\left(\frac{\hbar^2}{2m}\Delta + n_0U_0\right)\Phi(\mathbf{r},t) \\ &+\Phi(\mathbf{r},t)\int d\mathbf{r}'\, U(\mathbf{r}-\mathbf{r}')\,|\Phi(\mathbf{r}',t)|^2, \end{aligned} \tag{6}$$

where $U_0 = \int d\mathbf{r}'\, U(\mathbf{r}-\mathbf{r}')$. If one is interested only in the long-wave perturbations, and assumes that the characteristic spatial scale of the condensate wave function is much greater than the inter-particle interaction potential range, then the square of the wave function can be taken out of the integral in the right hand side of eq. (6). Thus, the GP equation takes the form

$$\begin{aligned} i\hbar\frac{\partial\Phi(\mathbf{r},t)}{\partial t} &= -\left(\frac{\hbar^2}{2m}\Delta + n_0U_0\right)\Phi(\mathbf{r},t) \\ &+U_0\Phi(\mathbf{r},t)\ |\Phi(\mathbf{r},t)|^2. \end{aligned} \tag{7}$$

This is the usual way to apply the GP equation. Small oscillations described by eq. (7) have the dispersion law

$$\omega^2 = \frac{U_0n_0k^2}{m} + \frac{\hbar^2k^4}{4m^2}, \tag{8}$$

where ω, k are the frequency and the wave number. The dispersion law (8) was derived by Bogolyubov [4] in a different approach with the use of the second quantization method. The excitation energy due to eq. (8) linearly depends on the wave number for small k and is monotone increasing with k, approaching the free particle dispersion law.

One should stress that spectrum (8) is valid only for the long-wave oscillations due to the approximation used for derivation of eq. (7).

Meanwhile, the derivation of the GP equation in form (6) does not assume that the excitations are of long-wave type. Thus, eq. (6) is also valid for description of the excitations which wave length is of order of the inter-particle interaction. We call further eq. (6) the non-local Gross-Pitaevskii equation (NGP). This equation can be applied in order to study the dispersion law of the collective excitations of a Bose-condensate in a short-wave domain of spectrum and to analyze the spatial behavior of the condensate wave function at the distances of order of the inter-atomic potential range.

HYDRODYNAMIC FORM OF NON-LOCAL GROSS-PITAEVSKII EQUATION

Equation (6) can be rewritten in the form of hydrodynamic equations, if the condensate wave function is taken as $\Phi = \eta e^{i\varphi}$, where φ, η are the phase and the absolute value of the condensate wave function and $\eta^2 = n$ is the particle number density in the condensate.

Taking the above into account, eq. (6) leads to

$$\frac{\partial n}{\partial t} + \nabla(n\mathbf{v}) = 0, \tag{9}$$

$$\begin{aligned} \frac{\hbar}{m}\frac{\partial\varphi}{\partial t} &= -\frac{\mathbf{v}^2}{2} + \frac{\hbar^2}{4m^2}\left(\frac{\Delta n}{n} - \frac{(\nabla n)^2}{2n^2}\right) \\ &+\frac{n_0U_0}{m} - \frac{1}{m}\int d\mathbf{r}'U(\mathbf{r}-\mathbf{r}')\,n(\mathbf{r}'), \end{aligned} \tag{10}$$

where the notion of superfluid velocity is introduced as

$$\mathbf{v} = (\hbar/m)\,\nabla\varphi. \tag{11}$$

Eq. (9) is the continuity equation, and eq. (10) is analogous to the Josephson equation in the theory of superconductivity. Taking a gradient of both sides of eq. (10) and taking into account the definition for the velocity, one gets

$$\frac{\partial\mathbf{v}}{\partial t} + (\mathbf{v}\nabla)\mathbf{v} = \nabla w, \tag{12}$$

where

$$w = \frac{\hbar^2}{2m}\frac{\Delta\eta}{\eta} - \frac{1}{m}\int d\mathbf{r}'U(\mathbf{r}-\mathbf{r}')\,n(\mathbf{r}').$$

In a linear approximation, the right hand side of eq. (12) can be expressed in form $\nabla w = -\nabla p/mn_0$, where the pressure and the density are related in a non-local way

$$p = -\frac{\hbar^2}{2}\eta_0\Delta\eta + n_0\int d\mathbf{r}'U(\mathbf{r}-\mathbf{r}')\,n(\mathbf{r}'), \tag{13}$$

Non-local relation between the pressure and the density was used in the phenomenological approach of [5] (neglecting the quantum term) in order to describe the rotons in the superfluid Helium. From our consideration it follows that the non-locality kernel, which relates the pressure and the density in a phenomenological approach, is determined by the inter-particle interaction potential.

SMALL OSCILLATION SPECTRUM WITH NON-LOCAL EFFECTS

The non-local GP equation leads to the following dispersion law for small oscillations

$$\omega^2 = \frac{U_k n_0 k^2}{m} + \frac{\hbar^2 k^4}{4m^2}, \tag{14}$$

which is different from (8) due to appearance of a potential Fourier component $U_k = \int d\mathbf{r}\, U(\mathbf{r})\, e^{ikr}$ (dependent on the wave vector) instead of U_0.

Usually, at short distances, a strong repulsion occurs, thus the interaction potential grows rapidly. It leads to a divergence of the Fourier component U_k. In order to overcome this obstacle one may suppose that the potential remains finite at short distances [6]. As the simplest case, we will use the „semitransparent sphere model“ potential

$$U(\mathbf{r}) = \begin{cases} U_m, & r \le a, \\ 0, & r > a, \end{cases} \tag{15}$$

and consider U_m and a as two independent parameters of the dimension of energy and length, correspondingly. In this case the Bose-condensate excitation spectrum depends on the only dimensionless parameter

$$\xi = \frac{\hbar^2}{8\pi\; n_0\; a^5\; m\; U_m}, \tag{16}$$

and can be written in form

$$\omega^2 = \omega_0^2 f(z), \tag{17}$$

where

$$\begin{aligned} f(z) &= z^{-1}(\sin z - z\cos z) + \xi z^4, && (18)\\ \omega_0^2 &= 4\pi\; n_0\; a\; U_m/m && (19)\\ z &= ka. && (20)\end{aligned}$$

Notice that the dimensionless parameter ξ (16) strongly depends on the potential range $\xi \propto a^{-5}$, rapidly growing with decreasing a.

The dependence of condensate excitation frequency on the wave number is shown in Fig.1. At parameter values $\xi > \xi_h = 3.4 \cdot 10^{-3}$, the dispersion curve does not have local minima and the energy monotone increases with wave number (upper curve in Fig.1).



Fig.1. Condensate excitation spectrum for the „semitransparent sphere model“ interaction; $\xi_h = 3.4 \cdot 10^{-3}$, $\xi_l = 0.92 \cdot 10^{-3}$

At $\xi_l < \xi < \xi_h$ the dispersion curve acquires a roton minimum (middle curve in Fig.1) and its depth increases with decreasing ξ. At the value $\xi = \xi_l = 0.92 \cdot 10^{-3}$, the excitation energy minimum equals to zero (lower curve in Fig.1); it means an instability of a many-particle system for given parameters of the interaction potential. Thus, in the range $\xi_l < \xi < \xi_h$ the excitation spectrum mimics that of superfluid Helium with a roton minimum [7].

The roton minimum position z_0 is determined by the equation $f'(z_0) = 0$. Dependence of z_0 on the parameter ξ is shown in Fig.2. With increasing ξ the roton minimum position gets shifted toward smaller wave numbers, changing from its maximum $z_{max} = 5.45$ (for $\xi_l = 0.92 \cdot 10^{-3}$) to minimum $z_{min} = 3.72$ (for $\xi_h = 3.4 \cdot 10^{-3}$). It follows that the excitation wave length λ, at the dispersion curve minimum lies in the range

$$1{,}15\,a < \lambda < 1{,}69\,a. \tag{21}$$

Thus one may conclude that the existence of a dispersion curve minimum is caused by the finiteness of the interparticle interaction range and is not connected with any specific superfluid properties of system. It is noteworthy that in Helium the repulsive range of the inter-atomic interaction potential may be estimated as $a = 2.56\,\text{Å}$ and the roton wave length is $\lambda_r = 3.3\,\text{Å}$ such that the relation $\lambda_r = 1.28\,a$ fits into the interval (21).

Dependence of the roton gap $\Delta = \omega_0\sqrt{f(z_0)}$ of the spectrum (17), (18) on the parameter ξ is monotone; Fig.3 shows that Δ/ω_0 is zero at $\xi = \xi_l$ and achieves its maximum $\Delta_{max}/\omega_0 = 1.17$ at $\xi = \xi_h$.



Fig.2. Roton minimum position z_0 as a function of parameter ξ



Fig.3 Roton gap versus parameter ξ.

ATTRACTION INFLUENCE ON THE SMALL OSCILLATION SPECTRUM

In the majority of the papers devoted to a non-ideal Bose-gas, as in [4], it is supposed that the inter-particle interaction is of repulsive type. It is of interest to investigate the influence of the inter-particle long-distance attraction on the elementary excitation spectrum.

Toward this end we calculate the spectrum for the interaction potential

$$U(\mathbf{r}) = \begin{cases} U_m, & r \le a, \\ -C/r^6, & r > a, \end{cases} \tag{22}$$

where C is positive. This potential accounts for a Van der Waals long-distance attraction of particles. The dispersion law for potential (22) can be expressed as

$$\omega^2 = \omega_0^2 \left(f(z) - \alpha g(z) \right), \tag{23}$$

with a dimensionless parameter $\alpha = C/U_m a^6$ and an auxiliary function

$$g(z) = \frac{z}{4}\left[\left(1-\frac{z^2}{6}\right)\sin z + \frac{z}{3}\left(1-\frac{z^2}{2}\right)\cos z + \frac{z^4}{6}\left(\frac{\pi}{2} - \mathrm{Si}(z)\right)\right], \tag{24}$$

($Si\,(z)$ stands for integral sine) describes the impact of the attractive part of the inter-particle potential on the elementary excitation spectrum. This function is shown in Fig.4. Function $g(z)$ is positive for wave numbers $0 < z < 2.45$ and negative for $2.45 < z < 5.40$. For $z > 5.40$ function $g\,(z)$ is positive and rises sharply.

One may notice that the interval $2.45 < z < 5.40$, where function (24) is negative, overlaps with the domain of the roton minimum existence. It follows that the attraction leads to the growth of the excitation energy in the domain of the roton minimum existence and smears this minimum. In the range of small wave numbers, and also for very large $z > 5.40$, attraction leads to lowering the excitation energy. Thus the long-distance atomic attraction results in smoothing of the dispersion curve and in narrowing the range of the existence of excitation spectrum, which has extrema (rotons and maxons). Attraction reduces the upper bound ξ_h and increases the lower one ξ_l, narrowing the range of the existence of a Landau like excitation spectrum $\delta\xi \equiv \xi_h - \xi_l$. Dependence of $\delta\xi$ for a spectrum with roton minimum on attraction strength $\alpha = C\left/U_m a^6\right.$, is shown in Fig.5.

Notice that experimental spectrum of quasi-particle excitations in the superfluid Helium at large momentum ($p > 3\text{Å}^{-1}$) increases slowly such that the derivative of $\omega(p)$ is close to zero at the boundary of the spectrum [8]. This behavior can be understood with help of a long-range attraction. Recall that function $g(z)$ is positive and increasing for $z > 5,4$. Thereby, due to (23), it opposes the dispersion curve growing at large momentum.

One should draw attention to smallness of the attraction part of the potential for Helium — it is the smallest among the rare gases. Namely, the depth of te potential well for Helium $\varepsilon = -10.5$K, for Neon $\varepsilon = -35.9$K, for Argon $\varepsilon = -121$K, for Krypton $\varepsilon = -173$K. It seems that due to this fact the maxon-roton part of the dispersion curve in liquid Helium is the most pronounced, not only in a superfluid phase, but in a normal phase as well. Notice a feature of the superfluid phase: the lifetime of the maxon-roton excitations are much larger than that in normal liquids.



Fig.4. Function $g(z)$ in the domain of z of physical interest.



Fig.5. Range $\delta\xi$ as a function of the interaction strength α.

CONCLUSIONS

It is shown that the Gross-Pitaevskii equation with account for the non-local effects, caused by the finiteness of the inter-particle interaction range, can describe the short-wave excitation spectrum, for which the wave length is of inter-particle interaction range order.

For a Bose-system the spectrum for arbitrary wave number is obtained; it has a shape of the Landau spectrum of a superfluid Helium. We estimated the range of parameters where such a spectrum exists. Computation shows the roton minimum in the dispersion curve is caused by the finiteness of the inter-particle interaction range; the wave length of this minimum is close to the repulsion range of the interaction. The influence of the long-distance inter-particle attraction on the dispersion curve is studied. It is shown that it results in smoothing of the excitation spectrum and in narrowing the range of the existence of a maxon-roton type curve. It is noticed that the attraction part of the potential for the Helium atoms is small in comparison with that for other atoms; it seems that this fact induces a pronounced maxon-roton part of the liquid Helium dispersion curve.

1. E.P. Gross, Nuovo Cimento **20**, 454 (1961).
2. L.P. Pitaevskii, JETP **40**, 646 (1961).
3. E.M. Lifshits, L.P Pitaevskii, Statistical Physics. Part.2 (Nauka, Moscow, 1978).
4. N.N. Bogolyubov, J. Phys. **9**, 23 (1947).
5. I.N. Adamenko , K.E. Nemchenko , I.V. Tanatarov, Phys. Rev. B **67**, 104513 (2003).
6. E.A. Pashitskii, Fiz. Nizk. Temp. **25**, N2, 115 (1999).
7. I.M. Khalatnikov, The theory of superfluidity (Nauka, Moscow, 1971).
8. H.R. Glyde, Phys. Rev. B **45**, 7321 (1992).

Received 18.09.09

КОРОТКОХВИЛЬОВІ ЗБУДЖЕННЯ В НЕЛОКАЛЬНІЙ МОДЕЛІ ГРОССА-ПІТАЄВСЬКОГО

А.П.Івашин, Ю.М.Полуектов

Резюме

Показано, що нелокальна форма рівняння Гросса-Пітаєвського дозволяє описати не тільки довгохвильові, але й короткохвильові збудження в системах з бозе-конденсатом. При певних значеннях параметрів, спектр збуджень бозе-конденсату нагадує спектр Ландау квазічастинкових збуджень у надплинному гелії з ротонним мінімумом. Довжина хвилі збуджень, при якій існує ротонний мінімум, близька до радіуса міжчастинкової взаємодії. Показано, що урахування міжчастинкового притягання звужує область існування спектра з ротонним мінімумом.

КОРОТКОВОЛНОВЫЕ ВОЗМУЩЕНИЯ В НЕЛОКАЛЬНОЙ МОДЕЛИ ГРОССА- ПИТАЕВСКОГО

А.П.Ивашин, Ю.М.Полуэктов

Резюме

Показано, что нелокальная форма уравнения Гросса-Питаевского позволяет описать не только длинноволновые, но и коротковолновые возбуждения в системах с бозе-конденсатом. При определенных значениях параметров, спектр возбуждений бозе-конденсата напоминает спектр Ландау квазичастичных возбуждений в сверхтекучем гелии с ротонным минимумом. Длина волны возбуждений, при которой существует ротонный минимум, близка к радиусу межчастичного взаимодействия. Показано, что учет межчастичного притяжения сужает область существования спектра с ротонным минимумом.